

University of Dundee

## Telomere length is an independent prognostic marker in MDS but not in de novo AML

Williams, Jenna; Heppel, Nicole H.; Britt-Compton, Bethan; Grimstead, Julia W.; Jones, Rhiannon E.; Tauro, Sudhir

*Published in:*  
British Journal of Haematology

*DOI:*  
[10.1111/bjh.14666](https://doi.org/10.1111/bjh.14666)

*Publication date:*  
2017

*Document Version*  
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

### *Citation for published version (APA):*

Williams, J., Heppel, N. H., Britt-Compton, B., Grimstead, J. W., Jones, R. E., Tauro, S., Bowen, D. T., Knapper, S., Groves, M., Hills, R. K., Pepper, C., Baird, D. M., & Fegan, C. (2017). Telomere length is an independent prognostic marker in MDS but not in *de novo* AML. *British Journal of Haematology*, 178(2), 240-249. <https://doi.org/10.1111/bjh.14666>

### General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



**University of Dundee**

## **Telomere length is an independent prognostic marker in MDS but not in de novo AML**

Fegan, Christopher; Williams, Jenna; Britt-Compton, Bethan; Heppel, Nicole; Grimstead, Julia; Jones, Rhiannon; Tauro, Sudhir; Bowen, David; Knapper, Steven; Groves, Michael; Hills, Robert; Pepper, Chris; Baird, Duncan

*Published in:*  
British Journal of Haematology

*Publication date:*  
2017

*Document Version*  
Accepted author manuscript

[Link to publication in Discovery Research Portal](#)

### *Citation for published version (APA):*

Fegan, C., Williams, J., Britt-Compton, B., Heppel, N., Grimstead, J., Jones, R., ... Baird, D. (2017). Telomere length is an independent prognostic marker in MDS but not in de novo AML. *British Journal of Haematology*.

### **General rights**

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



## Telomere length is an independent prognostic marker in MDS but not in de novo AML

Journal:	<i>British Journal of Haematology</i>
Manuscript ID	BJH-2016-01914.R1
Manuscript Type:	Ordinary Papers
Date Submitted by the Author:	10-Jan-2017
Complete List of Authors:	<p>Fegan, Christopher; University Hospital of Wales, Department of Haematology</p> <p>Williams, Jenna; Cardiff University School of Medicine, Cancer and Genetics</p> <p>Britt-Compton, Bethan; School of Medicine, Cancer &amp; Genetics</p> <p>Heppel, Nicole; School of Medicine, Cancer &amp; Genetics</p> <p>Grimstead, Julia; Cardiff University School of Medicine, Division of Cancer and Genetics</p> <p>Jones, Rhiannon; Cardiff University School of Medicine, Division of Cancer and Genetics</p> <p>Tauro, Sudhir; University of Dundee, Dundee Cancer Centre</p> <p>Bowen, David; Leeds General Infirmary, Department of Haematology</p> <p>Knapper, Steven; Cardiff University, Department of Haematology</p> <p>Groves, Michael; Ninewells Hospital, Haematology</p> <p>Hills, Robert; Cardiff University, Haematology</p> <p>Pepper, Chris; Cardiff University, Haematology</p> <p>Baird, Duncan; Cardiff University, School of Medicine</p>
Key Words:	AML, MDS, TELOMERE, TELOMERASE

Telomere length is prognostic in MDS but not AML

**Telomere length is an independent prognostic marker in MDS but not in de novo AML**

Jenna Williams,<sup>1</sup> Nicole H. Heppel,<sup>1</sup> Bethan Britt-Compton,<sup>1</sup> Julia W. Grimstead,<sup>1</sup> Rhiannon E. Jones,<sup>1</sup> Sudhir Tauro,<sup>2</sup> David T. Bowen,<sup>3</sup> Steven Knapper,<sup>1</sup> Michael Groves,<sup>2</sup> Robert K. Hills,<sup>1</sup> Chris Pepper,<sup>1</sup> Duncan M. Baird<sup>1</sup> and Chris Fegan.<sup>1</sup>

<sup>1</sup>Division of Cancer and Genetics, School of Medicine, Cardiff University, Heath Park, Cardiff CF14 4XN, UK

<sup>2</sup>Department of Haematology, Ninewells Hospital, Dundee, United Kingdom

<sup>3</sup>Department of Haematology, St James's Institute of Oncology, Leeds, United Kingdom

Corresponding author: Professor Chris Fegan, Division of Cancer and Genetics, School of Medicine, Cardiff University, Heath Park, Cardiff CF14 4XN, UK; e-mail: christopher.Fegan@wales.nhs.uk.

CF declares no conflict of interests other than co-authorship of a patent application based on some of this work; DMB declares no conflict of interests other than co-authorship of a patent application based on some of this work and CP declares no conflict of interests other than co-authorship of a patent application based on some of this work. All other authors have no conflicts of interest.

**Word count:** Summary 197, Main text 3195.

**Running title:** Telomere length is prognostic in MDS but not AML.

**Key words:** Myelodysplasia, AML, telomere, telomerase

<b>Number of figures</b>	3
<b>Number of Supplementary Figures</b>	3
<b>Number of tables</b>	1
<b>Number of references</b>	42

This is the peer reviewed version of the following article: 'Telomere length is an independent marker in MDS but not in de novo AML', British Journal of Haematology, which has been published in final form at <https://doi.org/10.1111/bjh.14666>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

Telomere length is prognostic in MDS but not AML

## 30 Summary

31 Telomere dysfunction is implicated in the generation of large-scale genomic rearrangements  
32 which drives progression to malignancy. In this study we used high-resolution single  
33 telomere length analysis (STELA) to examine the potential role of telomere dysfunction in 80  
34 Myelodysplasia (MDS) and 95 de novo Acute Myeloid Leukaemia (AML) patients. Despite  
35 the MDS cohort being older they had significantly longer telomeres than the AML cohort ( $P$   
36  $<.0001$ ) where telomere length was also significantly shorter in younger AML patients (age  
37  $<60$ ) ( $P = .02$ ) and in FLT3 ITD mutated AML patients ( $P = .03$ ). Using a previously determined  
38 telomere length threshold for telomere dysfunction (3.81kb) did not provide prognostic  
39 resolution in AML ( $HR = 0.68$ ,  $P = .2$ ). In contrast, the same length threshold was highly  
40 prognostic for overall survival in the MDS cohort ( $HR = 5.0$ ,  $P <.0001$ ). Furthermore, this  
41 telomere length threshold was an independent parameter in multivariate analysis when  
42 adjusted for age, gender, cytogenetic risk group, number of cytopenias and IPSS score ( $HR =$   
43  $2.27$ ,  $P <.0001$ ). Therefore, telomere length should be assessed in a larger prospective study  
44 to confirm its prognostic role in MDS with a view to integrating this variable into a revised  
45 IPSS.

Telomere length is prognostic in MDS but not AML

46     **Introduction**

47     The Myelodysplastic Syndromes (MDS) are a heterogeneous group of clonal haematopoietic  
48     disorders with varying survival and propensity to develop secondary acute myeloid  
49     leukaemia (sAML). In 1997 the International Prognostic Scoring System (IPSS) was based on  
50     the bone marrow blast count, 3 distinct cytogenetics risk groups and the number of  
51     cytopenias dividing patients into four IPSS subgroups(Greenberg, *et al* 1997). The IPSS – R  
52     (Revised) is an updated refinement of the IPSS which identifies five cytogenetic, three  
53     cytopenic and four blast count risk categories which combine into five overall prognostic  
54     subgroups(Greenberg, *et al* 2012, Vardiman, *et al* 2009). Many MDS patients have a normal  
55     karyotype but in recent years a very large number (up to 660) of molecular genetic defects  
56     have been identified encoding genes for cellular proteins including transcription factors e.g.  
57     *RUNX1*, epigenetic regulators and chromatin remodelling factors e.g. *TET2*, *DNMT3A*,  
58     *IDH1/2*, pre-RNA splicing factors e.g. *SF3B1*, receptor tyrosine kinase/ signalling molecules  
59     e.g. *NRAS*, *JAK2*, *NPM1*,*FLT3* and check point regulator *P53* some of which impact  
60     significantly on prognosis (Tothova, *et al* 2013, Walter, *et al* 2012). A recent MDS study  
61     showed that some of these somatic mutations were an independent prognostic marker  
62     compared to the IPSS (Bejar, *et al* 2011).

63     Recent studies have shown that the same genomic mutations seen in MDS/AML patients  
64     overlap with those identifiable in the normal adult population with increasing frequency  
65     with age (Genovese, *et al* 2014, Jaiswal, *et al* 2014, McKerrell, *et al* 2015, Xie, *et al* 2014).  
66     What drives these clones to further genomic instability and the development of MDS/AML is  
67     currently poorly understood. Telomeres are repetitive DNA sequences at the ends of  
68     chromosomes that shorten with each cell division. Critical loss of telomere length leads to  
69     chromosome end-end fusion and genomic instability resulting in large scale re-

Telomere length is prognostic in MDS but not AML

arrangements such as non-reciprocal translocations which are the hallmark of many tumour types including MDS, sAML and *de novo* AML (Jones, *et al* 2012). With ongoing cell division during life, the telomeres in normal individuals erode as a function of age at a rate of approximately 26 bp/year (Daniali, *et al* 2013). Several studies have suggested that MDS is associated with shorter telomeres leading to genomic instability and progression to sAML (Boulton, *et al* 1997, Chakraborty, *et al* 2009, Sieglöva, *et al* 2004, Townsley, *et al* 2014, Young 2010). Recently, using a modified Q-FISH based method, Gadji *et al* proposed that telomere dysfunction underpins the chromosomal changes associated with MDS progression to AML and *de novo* AML (Gadji, *et al* 2012). Whilst in a mouse model it was shown that telomere dysfunction induced the same types of DNA damage that drives classical MDS phenotypes including *SF3B1* and *DNMT3A* resulting in differentiation changes in myeloid precursors (Colla, *et al* 2015).

We previously developed a high-resolution technique to determine telomere length, Single Telomere Length Analysis (STELA), which is unique in its ability to detect telomere lengths from single chromosomes within the length ranges at which telomere fusions can occur (Lin, *et al* 2010, Lin, *et al* 2014). Using STELA we have shown that some chronic lymphocytic leukaemia (CLL) patients display extreme telomere shortening and fusion consistent with the onset of a telomere-driven crisis that can drive the formation of large-scale genome rearrangements (Lin, *et al* 2010). Telomeres in these ranges cannot be readily detected with the other methodologies such as QPCR, Southern blot or Q-FISH previously used in MDS/AML studies (Baird, *et al* 2003, Britt-Compton, *et al* 2012, Chakraborty, *et al* 2009, Lin, *et al* 2010). We showed that in CLL patients, telomere fusions only occurred when telomere length was  $\leq 3.81$  kb and importantly we show that telomere erosion to within these length

Telomere length is prognostic in MDS but not AML

93 ranges precedes clinical progression. Indeed high-resolution telomere length analysis using  
94 STELA, together with the stratification of patients based on the telomere length thresholds  
95 at which fusion occurs, provided an independent high-resolution marker of prognosis even  
96 in patients with early-stage CLL (Lin, *et al* 2014).

97 In this study we used STELA to assess if telomere erosion is an important pathogenic  
98 mechanism driving prognosis MDS and AML.

99 **Methods**

100 **Patients, samples and cell separation**

101 This study was undertaken at the University Hospital of Wales (UHW), Cardiff. All of the  
102 unselected diagnostic patient bone marrow samples: 80 MDS which consisted of 37 samples  
103 from Dundee (archived from 1997 – 2005) and 43 from Cardiff (archived between 1985 and  
104 2008) including 7 with a proven and documented history of MDS prior to progression to  
105 AML (blasts >20%), and 95 de novo AML from Cardiff (archived between 2003 and 2012)  
106 were obtained following written informed consent (Table 1). We deliberately chose to use  
107 stored but well annotated archival samples so as to be able to assess the potential impact of  
108 telomere length on survival in all patient groups but especially low risk MDS patients. All  
109 MDS patients were treated with best supportive care with none receiving azacitidine prior  
110 to this analysis. All morphological, immunophenotypic, cytogenetic and molecular data  
111 were collected from Cardiff and Dundee Haematology Departments for their respective  
112 patients but all telomere length analyses on the bone marrow mononuclear cells were  
113 undertaken within the Division of Cancer and Genetics, Cardiff University. **Bone marrow**  
114 **mononuclear cells were collected in ethylenediaminetetraacetic acid and isolated by**  
115 **density centrifugation using Ficoll-Hypaque (Invitrogen) which resulted in <3%**



Telomere length is prognostic in MDS but not AML

lymphocytes contamination. Cells were stored at  $-20^{\circ}\text{C}$  as dry pellets before DNA extraction.

DNA extraction, single telomere length analysis and telomerase assays.

Telomere length was determined using XpYp STELA as previously described (Lin, *et al* 2010, Roger, *et al* 2013). Briefly, DNA was extracted using proteinase K, RNase A, phenol/chloroform protocols and quantified by Hoechst 33258 fluorometry (Bio-Rad) before dilution to  $10\text{ ng}/\mu\text{L}$  in  $10\text{mM}$  Tris-HCl, pH 7.5. A total of  $10\text{ ng}$  of DNA was further diluted to  $250\text{ pg}/\mu\text{L}$  in a volume of  $40\text{ }\mu\text{L}$  containing  $1\mu\text{M}$  Telorette2 linker and  $1\text{mM}$  Tris-HCl, pH 7.5. Multiple polymerase chain reactions (PCRs; typically 6 reactions per sample) were carried out for each test DNA in  $10\text{-}\mu\text{L}$  volumes  $250\text{ pg}$  of DNA,  $0.5\mu\text{M}$  of the telomere-adjacent and Teltail primers,  $75\text{mM}$  Tris-HCl, pH 8.8,  $20\text{mM}$   $(\text{NH}_4)_2\text{SO}_4$ ,  $0.01\%$  Tween-20,  $1.5\text{mM}$   $\text{MgCl}_2$ , and  $0.5\text{ U}$  of a  $10:1$  mixture of Taq (ABGene) and Pwo polymerase (Roche Molecular Biochemicals). The reactions were cycled with an MJ PTC-225 thermocycler (MJ Research). The DNA fragments were resolved by  $0.5\%$  Tris acetate ethylenediaminetetraacetic acid agarose gel electrophoresis, and detected by Southern blot hybridization with random-primed  $\alpha\text{-}^{33}\text{P}$ -labelled (GE Healthcare) TTAGGG repeat probe together with probes to detect the  $1\text{-kb}$  (Stratagene) and  $2.5\text{-kb}$  (Bio-Rad) molecular weight markers. The hybridized fragments were detected by phosphorimaging with a Molecular Dynamics Storm 860 phosphorimager (GE Healthcare). The molecular weights of the DNA fragments were calculated using the Phoretix 1D quantifier (Nonlinear Dynamics). Telomerase assays were undertaken using the TRAPeze XL Telomerase detection kit (Chemicon International, Billerica, MA) as previously described (Lin, *et al* 2010, Roger, *et al* 2013).

Telomere length is prognostic in MDS but not AML

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

139     **Statistical methods**

140     Statistical analysis was undertaken by the Haematology Clinical Trials Unit, Cardiff University  
141     using SAS version 9.4 and GraphPad Prism 6. Spearman’s correlation was used for  
142     correlations between baseline values; Mann-Whitney and Wilcoxon matched-paired  
143     nonparametric tests were used for comparisons between groups. Mean telomere length  
144     was assessed between diagnoses using the Wilcoxon rank sum test. Paired data were  
145     compared using the Wilcoxon signed rank test. **In the MDS cohort patients were assessed**  
146     **using the 1997 IPSS criteria for the 3 cytogenetic risk groups, blast count, number of**  
147     **cytopenias (Hb <100g/l, neutrophils <1.5x10<sup>9</sup>/l and platelets <100x10<sup>9</sup>/l), IPSS score, age**  
148     **and sex whereas the AML cohort was assessed for cytogenetic risk group and FLT3 and**  
149     **NPM mutations (Falini, *et al* 2005, Kottaridis, *et al* 2001, Townsley, *et al* 2014).** Survival  
150     was assessed using the Kaplan-Meier method, and compared using Cox proportional  
151     hazards regression, with model building carried out using forward selection with significance  
152     set at P = .05.

153     **Results**

154     **Telomere length MDS and AML patients.**

155     An MDS patient’s bone marrow will typically display a variable range of immature and more  
156     differentiated cells which may or may not be derived from the malignant stem cells. We  
157     therefore compared the telomere length of the first 20 individual MDS patients CD34+ and  
158     CD34- bone marrow cells, but found no significant difference in telomere length between  
159     the two fractions (P = .08; Supplementary Figure 1A). We therefore proceeded without cell

Telomere length is prognostic in MDS but not AML

selection in subsequent MDS patient bone marrow sample analyses and simply used bone marrow mononuclear cell pellets.

Despite the fact that the MDS patients (median age 68 years, range 21-86) were older than AML patients (median age 56 years, range 17-80) telomere length was significantly longer in the MDS cohort compared to the AML cohort ( $P < .0001$ ; Figure 1A). Although MDS samples showed a modest reduction in telomere length with increasing age at diagnosis, there was no significant correlation ( $\rho^2 = -.0212$ ,  $P = .2$ ; Figure 1B) whereas in the AML samples a positive correlation was observed ( $\rho^2 = .0890$ ,  $P = .003$ ; Figure 1C). This finding is in contrast to what occurs during normal ageing; samples derived from older AML patients (age  $>60$ ) had significantly longer telomeres than younger AML patients ( $P = .02$ ; Figure 1D). Indeed, 22/26 (84.6%) of AML patients age  $<50$  years had telomere lengths within the range at which they can become dysfunctional or 'fusogenic' ( $\leq 3.81$  kb), which we previously described in CLL (Lin, *et al* 2014).

173

#### 174 **Telomere length, blast count, cytogenetics, cytopenias and IPSS.**

We next analysed the MDS cohort to assess any possible correlations between telomere length and age, gender, blast count, number of cytopenias, cytogenetic risk group and IPSS sub-group. Shorter telomere length was associated with male gender ( $P = .01$ ; Supplementary Figure 1B) and increased number of cytopenias ( $P = .003$ ; Figure 2A). None of the other parameters were significantly associated with telomere length. Consistent with previous reports we found a significant association between the number of cytopenias and overall survival ( $P < .0001$ ; Supplementary Figure 2A). Furthermore, patients with high risk

Telomere length is prognostic in MDS but not AML

1  
2  
3 182 cytogenetic abnormalities tended to have worse survival although this did not reach  
4  
5 183 statistical significance ( $P = .12$ ; Supplementary Figure 2B). Interestingly we found no simple  
6  
7  
8 184 association between telomere length and the three IPSS cytogenetic risk groups ( $P = .6$  for  
9  
10 185 trend; Figure 2B), the four IPSS subgroups ( $\rho^2 = .14$  for correlation; Figure 2C) or bone  
11  
12 186 marrow blast percentage ( $\rho^2 = -0.22$ ,  $P = .0503$  Figure 2D).

13  
14  
15  
16 187 In the de novo AML cohort we assessed the correlation between telomere length and  
17  
18 188 gender, age, presenting WBC, performance status, whether primary or secondary AML and  
19  
20 189 NPM1 and FLT3 mutation status. There was no association between NPM1 mutated patients  
21  
22 190 and telomere length (data not shown), but significantly shorter telomeres were found in the  
23  
24 191 FLT3 ITD mutated group when compared to FLT3 wild type AML patients ( $P = .03$ ). In  
25  
26 192 contrast, there was a trend towards longer telomeres in the FLT3 TKD mutated group  
27  
28 193 compared to ITD mutated AML ( $P = .12$ ; Figure 2E).

29  
30  
31  
32  
33 194

34  
35  
36 195 **Telomerase activity in MDS and AML patients.**

37  
38  
39 196 Given the different telomere length characteristics of the MDS and AML cohorts, we  
40  
41 197 investigated whether this may reflect differences in telomerase activity, an enzyme  
42  
43 198 responsible for extending shortened telomeres. Telomerase activity was analysed in a  
44  
45 199 subset of CD34+ AML samples ( $n = 12$ ) and in purified CD34+ cells from MDS patients ( $n =$   
46  
47 200 20). Telomerase activity was significantly higher in the AML samples when compared with  
48  
49 201 CD34+ MDS cells ( $P = .0002$ ; Figure 2F).

50  
51  
52  
53  
54 202

55  
56  
57 203 **Telomere length and survival in MDS and AML.**

Telomere length is prognostic in MDS but not AML

204 We next assessed the impact of telomere length on overall survival in the MDS and AML  
205 cohorts. Segregation of the two cohorts according to whether their mean telomere length  
206 was above or below the upper limit of telomere dysfunction (3.81 kb) revealed no  
207 difference in survival in the AML patient group (HR = 0.68 (0.37-1.9),  $P = .2$ ; Figure 3A). In  
208 contrast, bifurcation of the MDS cohort using this threshold telomere length demonstrated  
209 that patients with a median telomere length  $\leq 3.81$  kb had significantly worse survival (HR =  
210 5.0 (2.7-10.0),  $P < .0001$ ; Figure 3B). The impact of telomere length in these two disease  
211 settings was confirmed using telomere length quartile analysis. There was no correlation  
212 between telomere length and overall survival or relapse-free survival in the AML cohort ( $P =$   
213  $.5$ ,  $P = .09$ ; Supplementary Figures 3A and 3B). However, similar quartile analysis in the MDS  
214 cohort clearly showed that patients in the lowest two quartiles had worse survival with all  
215 patients in the lowest quartile alive at 3 years (Supplementary Figure 3C).

#### 217 **Telomere length is an independent prognostic variable in MDS.**

218 Finally, we performed multivariate analysis in the MDS cohort using a forward selection  
219 model that included age, IPSS, gender, cytogenetic risk group, number of cytopenias and  
220 telomere length. Short telomere length was identified as the most significant independent  
221 marker of overall survival (HR = 2.27 (1.45-3.57),  $P < .0001$ ). When entering short telomere  
222 length into the model the following parameters retained independent prognostic value: high  
223 IPSS (HR = 1.2 (0.64-2.27), increased number of cytopenias (HR = 1.60 (1.09-2.35),  $P = .007$ ),  
224 older age (HR per year 1.03 (1.01-1.07),  $P = .05$ ) and male gender (HR = 2.70 (1.20-6.10),  $P =$   
225  $.01$ ).

Telomere length is prognostic in MDS but not AML

Discussion

We previously showed that a proportion of chronic lymphocytic leukaemia (CLL) patients display extreme telomere erosion and fusion consistent with a telomere-driven crisis. Importantly, this was not simply a function of advanced stage disease but was detected in a subset of early stage patients prior to clinical progression (Lin, *et al* 2010). Subsequently we showed that defining the specific telomere length threshold at which telomere fusion occurred was a powerful way to risk-stratify CLL patients even those with early-stage disease (Lin, *et al* 2014). Previous studies had suggested a possible role for telomere dysfunction in both MDS and AML, so here we investigated the relationship between telomere length, disease progression and clinical outcome in MDS and AML using the high-resolution STELA technique.

We demonstrated that telomere length was highly predictive of disease outcome in MDS. In contrast, we found no evidence that telomere length influenced the survival of de novo AML patients. It should be noted, however, that we demonstrated shorter telomere lengths in the FLT3-ITD mutated group, a group with a well-established poorer prognosis. These results are similar to those of Aalbers *et al* who also showed shorter telomeres in patients with FLT3-ITD mutations but not NPM1 mutated patients (Aalbers, *et al* 2013). One possible explanation for the different telomere length characteristics in MDS and AML may be the differential expression of telomerase found in these two conditions. Unlike MDS cells, AML cells showed evidence of upregulated telomerase activity, which could prevent replicative

Telomere length is prognostic in MDS but not AML

249 senescence and allow unlimited proliferation during leukaemogenesis (Engelhardt, *et al*  
250 2004, Shay and Wright 2011). MDS is characterised by clonal expansion and hypercellularity  
251 but increased apoptosis leading to cytopenias (Greenberg 1999, Greenberg 1998, Parker, *et*  
252 *al* 1998, Parker and Mufti 2000). The reason why telomerase is not upregulated in MDS is  
253 unknown, but to date studies have failed to show any responsible acquired telomerase-  
254 regulated genetic abnormalities in this patient group, although there are several germline  
255 mutations described in TERC and TERT predisposing to MDS/AML (Ballew and Savage 2013,  
256 Calado, *et al* 2009, Kirwan, *et al* 2009, Ohyashiki, *et al* 1999, Yamaguchi, *et al* 2003).

257 Our study showed an association between telomere length and the number of cytopenias in  
258 MDS patients. It seems possible that the lack of telomerase activity observed in MDS leads  
259 to unhindered cell senescence and/or apoptosis. Also the high proliferative and apoptotic  
260 rates seen in MDS bone marrow produces increased ineffective haematopoiesis and more  
261 profound cytopenias (Raza, *et al* 1997a, Raza, *et al* 1997b). These results are similar to  
262 those of Sieglöva *et al* who showed MDS patients with shorter telomeres were more likely  
263 to progress to AML (Sieglöva, *et al* 2004). Perhaps surprisingly, there was no association  
264 between telomere length and cytogenetic risk group, blast count and IPSS score. This lack of  
265 association meant that telomere length was an independent marker of outcome in MDS and  
266 in the multivariate forward selection model we employed. Indeed, it was more prognostic  
267 than blast count, cytogenetics, number of cytopenias or IPSS in this context.

268 Several recent studies have shown that clonal haematopoiesis is almost a “normal” part of  
269 ageing with recent reports showing 0.8%, 11% and 19.5 % of normal individuals aged <60,  
270 >80 and >90 years respectively having demonstrable clonal haematopoiesis – so called age-  
271 related clonal haematopoiesis. These include the acquisition of many genetic lesions

Telomere length is prognostic in MDS but not AML

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

272 associated with the development of MDS/AML including DNMT3, IDH1, IDH2, NRAS, KRAS,  
273 JAK 2, SF3B1 and SRSF2 mutations (Jaiswal, *et al* 2014, McKerrell, *et al* 2015, Xie, *et al* 2014).  
274 Three of our observations in this study were that the telomere length increases with age in  
275 AML patients, that AML patients older than 60 years having significantly long telomeres  
276 than those age <60 years and that the AML patient cohort had significantly shorter  
277 telomeres than the MDS patient cohort despite being younger. One possible explanation for  
278 these data is that older de novo AML patients may have developed secondary AML despite  
279 the absence of a documented history of prior MDS or clonal haematopoiesis. This would be  
280 in keeping with the recent identification of increased genomic instability in ageing “normal”  
281 people with many identical genomic abnormalities seen in elderly AML patients (McKerrell,  
282 *et al* 2015). The acquisition of the various differing leukaemia-associated genomic  
283 abnormalities are age dependent with for example the recurrent point mutations affecting  
284 spliceosome genes SF3B1 and SRSF2 associated with clonal haematopoiesis in individuals  
285 aged 70 years or over, but not in younger people (McKerrell and Vassiliou 2015,  
286 Papaemmanuil, *et al* 2013). An alternative, but not exclusive hypothesis, is that AML in  
287 younger patients tends be more progenitor-type AML (Core binding factor, NPM1-mutated,  
288 FLT3-ITD-mutated) and more proliferative leading to shorter telomeres. Further studies are  
289 required to assess the relationship of shorter telomeres with age-related clonal  
290 haematopoiesis and MDS and AML development.

291 The findings in MDS are in keeping with our previous data in CLL and breast cancer where  
292 we demonstrated the utility of our telomere-length threshold in providing powerful  
293 independent prognostic information (Lin, *et al* 2014, Roger, *et al* 2013). Taken together  
294 these data point to a common mechanism that is present in diverse tumour types, by which



Telomere length is prognostic in MDS but not AML

295 the presence of short dysfunctional telomeres can drive genomic instability and clonal  
296 evolution leading to poor clinical outcomes.

297 Our study does have several limitations in that it consists of relatively small cohorts of MDS  
298 and AML patients and was deliberately retrospective so as to be able to assess the impact of  
299 telomere length on survival especially in low risk MDS patients, all of whom were treated  
300 with supportive care only. Finally our observation that telomere length is independently  
301 prognostic in MDS indicates that consideration should be given to a much larger prospective  
302 study assessing the potential role of telomeres in the prognostication of MDS. This would  
303 also facilitate the assessment of telomere length analysis as a potential predictor of  
304 response to newer therapies such as azacitidine and perhaps its ultimate incorporation into  
305 a revised IPSS.

#### 306 **Acknowledgements.**

307 We acknowledge the Tayside Tissue Bank for supplying some of the MDS samples and in  
308 particular Norene Keenan, and Ann Hyslop.

309 Performed the research – JW, MHH, BB-C, JWG, REJ, DMB

310 Designed the research study – CF, CP, DMB

311 Contributed essential reagents or tools –ST, MG, DTB, SK

312 Analysed the data – CP, DMB, RKH

313 Wrote the paper – CP, DMB, CF

314 This work was funded by Cancer Research UK and Bloodwise grants 12049 and 13033.

Telomere length is prognostic in MDS but not AML

References

Aalbers, A.M., Calado, R.T., Young, N.S., Zwaan, C.M., Wu, C., Kajigaya, S., Coenen, E.A., Baruchel, A., Geleijns, K., de Haas, V., Kaspers, G.J., Kuijpers, T.W., Reinhardt, D., Trka, J., Zimmermann, M., Pieters, R., van der Velden, V.H. & van den Heuvel-Eibrink, M.M. (2013) Telomere length and telomerase complex mutations in pediatric acute myeloid leukemia. *Leukemia*, **27**, 1786-1789.

Baird, D.M., Rowson, J., Wynford-Thomas, D. & Kipling, D. (2003) Extensive allelic variation and ultrashort telomeres in senescent human cells. *Nature genetics*, **33**, 203-207.

Ballew, B.J. & Savage, S.A. (2013) Updates on the biology and management of dyskeratosis congenita and related telomere biology disorders. *Expert review of hematology*, **6**, 327-337.

Bejar, R., Stevenson, K., Abdel-Wahab, O., Galili, N., Nilsson, B., Garcia-Manero, G., Kantarjian, H., Raza, A., Levine, R.L., Neuberg, D. & Ebert, B.L. (2011) Clinical effect of point mutations in myelodysplastic syndromes. *The New England journal of medicine*, **364**, 2496-2506.

Boultonwood, J., Fidler, C., Kusec, R., Rack, K., Elliott, P.J., Atoyebi, O., Chapman, R., Oscier, D.G. & Wainscoat, J.S. (1997) Telomere length in myelodysplastic syndromes. *American journal of hematology*, **56**, 266-271.

Britt-Compton, B., Lin, T.T., Ahmed, G., Weston, V., Jones, R.E., Fegan, C., Oscier, D.G., Stankovic, T., Pepper, C. & Baird, D.M. (2012) Extreme telomere erosion in ATM-mutated and 11q-deleted CLL patients is independent of disease stage. *Leukemia : official journal of the Leukemia Society of America, Leukemia Research Fund, U.K.*, **26**, 826-830.

Calado, R.T., Regal, J.A., Hills, M., Yewdell, W.T., Dalmazzo, L.F., Zago, M.A., Lansdorp, P.M., Hogge, D., Chanock, S.J., Estey, E.H., Falcao, R.P. & Young, N.S. (2009) Constitutional hypomorphic telomerase mutations in patients with acute myeloid leukemia. *Proceedings of the National Academy of Sciences of the United States of America*, **106**, 1187-1192.

Chakraborty, S., Sun, C.L., Francisco, L., Sabado, M., Li, L., Chang, K.L., Forman, S., Bhatia, S. & Bhatia, R. (2009) Accelerated telomere shortening precedes development of therapy-related myelodysplasia or acute myelogenous leukemia after autologous transplantation for lymphoma. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*, **27**, 791-798.

Colla, S., Ong, D.S., Ogoti, Y., Marchesini, M., Mistry, N.A., Clise-Dwyer, K., Ang, S.A., Storti, P., Viale, A., Giuliani, N., Ruisaard, K., Ganan Gomez, I., Bristow, C.A., Estecio, M., Weksberg, D.C., Ho, Y.W., Hu, B., Genovese, G., Pettazzoni, P., Multani, A.S., Jiang, S., Hua, S., Ryan, M.C., Carugo, A., Nezi, L., Wei, Y., Yang, H., D'Anca, M., Zhang, L., Gaddis, S., Gong, T., Horner, J.W., Heffernan, T.P., Jones, P., Cooper, L.J., Liang, H., Kantarjian, H., Wang, Y.A., Chin, L., Bueso-Ramos, C., Garcia-Manero, G. & DePinho, R.A. (2015) Telomere dysfunction drives aberrant hematopoietic differentiation and myelodysplastic syndrome. *Cancer Cell*, **27**, 644-657.

Daniali, L., Benetos, A., Susser, E., Kark, J.D., Labat, C., Kimura, M., Desai, K., Granick, M. & Aviv, A. (2013) Telomeres shorten at equivalent rates in somatic tissues of adults. *Nat Commun*, **4**, 1597.

Engelhardt, M., Wasch, R. & Guo, Y. (2004) Telomeres and telomerase in normal and leukemic hematopoietic cells. *Leukemia research*, **28**, 1001-1004.

Falini, B., Mecucci, C., Tiacci, E., Alcalay, M., Rosati, R., Pasqualucci, L., La Starza, R., Diverio, D., Colombo, E., Santucci, A., Bigerna, B., Pacini, R., Pucciarini, A., Liso, A., Vignetti, M., Fazi, P., Meani, N., Pettirossi, V., Saglio, G., Mandelli, F., Lo-Coco, F., Pelicci, P.G. & Martelli, M.F. (2005) Cytoplasmic nucleophosmin in acute myelogenous leukemia with a normal karyotype. *The New England journal of medicine*, **352**, 254-266.

Gadji, M., Adebayo Awe, J., Rodrigues, P., Kumar, R., Houston, D.S., Klewes, L., Dieye, T.N., Rego, E.M., Passetto, R.F., de Oliveira, F.M. & Mai, S. (2012) Profiling three-dimensional nuclear

Telomere length is prognostic in MDS but not AML

- telomeric architecture of myelodysplastic syndromes and acute myeloid leukemia defines patient subgroups. *Clinical cancer research : an official journal of the American Association for Cancer Research*, **18**, 3293-3304.
- Genovese, G., Kahler, A.K., Handsaker, R.E., Lindberg, J., Rose, S.A., Bakhoum, S.F., Chambert, K., Mick, E., Neale, B.M., Fromer, M., Purcell, S.M., Svantesson, O., Landen, M., Hoglund, M., Lehmann, S., Gabriel, S.B., Moran, J.L., Lander, E.S., Sullivan, P.F., Sklar, P., Gronberg, H., Hultman, C.M. & McCarroll, S.A. (2014) Clonal hematopoiesis and blood-cancer risk inferred from blood DNA sequence. *The New England journal of medicine*, **371**, 2477-2487.
- Greenberg, P. (1999) Apoptosis and its role in myelodysplastic syndrome. *Leukemia research*, **23**, 855.
- Greenberg, P., Cox, C., LeBeau, M.M., Fenaux, P., Morel, P., Sanz, G., Sanz, M., Vallespi, T., Hamblin, T., Oscier, D., Ohyashiki, K., Toyama, K., Aul, C., Mufti, G. & Bennett, J. (1997) International scoring system for evaluating prognosis in myelodysplastic syndromes. *Blood*, **89**, 2079-2088.
- Greenberg, P.L. (1998) Apoptosis and its role in the myelodysplastic syndromes: implications for disease natural history and treatment. *Leukemia research*, **22**, 1123-1136.
- Greenberg, P.L., Tuechler, H., Schanz, J., Sanz, G., Garcia-Manero, G., Sole, F., Bennett, J.M., Bowen, D., Fenaux, P., Dreyfus, F., Kantarjian, H., Kuendgen, A., Levis, A., Malcovati, L., Cazzola, M., Cermak, J., Fonatsch, C., Le Beau, M.M., Slovak, M.L., Krieger, O., Luebbert, M., Maciejewski, J., Magalhaes, S.M., Miyazaki, Y., Pfeilstocker, M., Sekeres, M., Sperr, W.R., Stauder, R., Tauro, S., Valent, P., Vallespi, T., van de Loosdrecht, A.A., Germing, U. & Haase, D. (2012) Revised international prognostic scoring system for myelodysplastic syndromes. *Blood*, **120**, 2454-2465.
- Jaiswal, S., Fontanillas, P., Flannick, J., Manning, A., Grauman, P.V., Mar, B.G., Lindsley, R.C., Mermel, C.H., Burt, N., Chavez, A., Higgins, J.M., Moltchanov, V., Kuo, F.C., Kluk, M.J., Henderson, B., Kinnunen, L., Koistinen, H.A., Ladenvall, C., Getz, G., Correa, A., Banahan, B.F., Gabriel, S., Kathiresan, S., Stringham, H.M., McCarthy, M.I., Boehnke, M., Tuomilehto, J., Haiman, C., Groop, L., Atzmon, G., Wilson, J.G., Neuberg, D., Altshuler, D. & Ebert, B.L. (2014) Age-related clonal hematopoiesis associated with adverse outcomes. *The New England journal of medicine*, **371**, 2488-2498.
- Jones, C.H., Pepper, C. & Baird, D.M. (2012) Telomere dysfunction and its role in haematological cancer. *British journal of haematology*, **156**, 573-587.
- Kirwan, M., Beswick, R., Vulliamy, T., Nathwani, A.C., Walne, A.J., Casimir, C. & Dokal, I. (2009) Exogenous TERC alone can enhance proliferative potential, telomerase activity and telomere length in lymphocytes from dyskeratosis congenita patients. *British journal of haematology*, **144**, 771-781.
- Kottaridis, P.D., Gale, R.E., Frew, M.E., Harrison, G., Langabeer, S.E., Belton, A.A., Walker, H., Wheatley, K., Bowen, D.T., Burnett, A.K., Goldstone, A.H. & Linch, D.C. (2001) The presence of a FLT3 internal tandem duplication in patients with acute myeloid leukemia (AML) adds important prognostic information to cytogenetic risk group and response to the first cycle of chemotherapy: analysis of 854 patients from the United Kingdom Medical Research Council AML 10 and 12 trials. *Blood*, **98**, 1752-1759.
- Lin, T.T., Letsolo, B.T., Jones, R.E., Rowson, J., Pratt, G., Hewamana, S., Fegan, C., Pepper, C. & Baird, D.M. (2010) Telomere dysfunction and fusion during the progression of chronic lymphocytic leukemia: evidence for a telomere crisis. *Blood*, **116**, 1899-1907.
- Lin, T.T., Norris, K., Heppel, N.H., Pratt, G., Allan, J.M., Allsup, D.J., Bailey, J., Cawkwell, L., Hills, R., Grimstead, J.W., Jones, R.E., Britt-Compton, B., Fegan, C., Baird, D.M. & Pepper, C. (2014) Telomere dysfunction accurately predicts clinical outcome in chronic lymphocytic leukaemia, even in patients with early stage disease. *British journal of haematology*, **167**, 214-223.
- McKerrell, T., Park, N., Moreno, T., Grove, C.S., Ponstingl, H., Stephens, J., Crawley, C., Craig, J., Scott, M.A., Hodgkinson, C., Baxter, J., Rad, R., Forsyth, D.R., Quail, M.A., Zeggini, E., Ouwehand, W.,

Telomere length is prognostic in MDS but not AML

Varela, I. & Vassiliou, G.S. (2015) Leukemia-associated somatic mutations drive distinct patterns of age-related clonal hemopoiesis. *Cell reports*, **10**, 1239-1245.

McKerrell, T. & Vassiliou, G.S. (2015) Aging as a driver of leukemogenesis. *Sci Transl Med*, **7**, 306fs338.

Ohyashiki, J.H., Iwama, H., Yahata, N., Ando, K., Hayashi, S., Shay, J.W. & Ohyashiki, K. (1999) Telomere stability is frequently impaired in high-risk groups of patients with myelodysplastic syndromes. *Clinical cancer research : an official journal of the American Association for Cancer Research*, **5**, 1155-1160.

Papaemmanuil, E., Gerstung, M., Malcovati, L., Tauro, S., Gundem, G., Van Loo, P., Yoon, C.J., Ellis, P., Wedge, D.C., Pellagatti, A., Shlien, A., Groves, M.J., Forbes, S.A., Raine, K., Hinton, J., Mudie, L.J., McLaren, S., Hardy, C., Latimer, C., Della Porta, M.G., O'Meara, S., Ambaglio, I., Galli, A., Butler, A.P., Walldin, G., Teague, J.W., Quek, L., Sternberg, A., Gambacorti-Passerini, C., Cross, N.C., Green, A.R., Boultonwood, J., Vyas, P., Hellstrom-Lindberg, E., Bowen, D., Cazzola, M., Stratton, M.R., Campbell, P.J. & Chronic Myeloid Disorders Working Group of the International Cancer Genome, C. (2013) Clinical and biological implications of driver mutations in myelodysplastic syndromes. *Blood*, **122**, 3616-3627; quiz 3699.

Parker, J.E., Fishlock, K.L., Mijovic, A., Czepulkowski, B., Pagliuca, A. & Mufti, G.J. (1998) 'Low-risk' myelodysplastic syndrome is associated with excessive apoptosis and an increased ratio of pro- versus anti-apoptotic bcl-2-related proteins. *British journal of haematology*, **103**, 1075-1082.

Parker, J.E. & Mufti, G.J. (2000) Excessive apoptosis in low risk myelodysplastic syndromes (MDS). *Leukemia & lymphoma*, **40**, 1-24.

Raza, A., Alvi, S., Borok, R.Z., Span, L., Parcharidou, A., Alston, D., Rifkin, S., Robin, E., Shah, R. & Gregory, S.A. (1997a) Excessive proliferation matched by excessive apoptosis in myelodysplastic syndromes: the cause-effect relationship. *Leukemia & lymphoma*, **27**, 111-118.

Raza, A., Alvi, S., Broady-Robinson, L., Showel, M., Cartlidge, J., Mundle, S.D., Shetty, V.T., Borok, R.Z., Dar, S.E., Chopra, H.K., Span, L., Parcharidou, A., Hines, C., Gezer, S., Venugopal, P., Loew, J., Showel, J., Alston, D., Hernandez, B., Rifkin, S., Robin, E., Shah, R. & Gregory, S.A. (1997b) Cell cycle kinetic studies in 68 patients with myelodysplastic syndromes following intravenous iodo- and/or bromodeoxyuridine. *Experimental hematology*, **25**, 530-535.

Roger, L., Jones, R.E., Heppel, N.H., Williams, G.T., Sampson, J.R. & Baird, D.M. (2013) Extensive telomere erosion in the initiation of colorectal adenomas and its association with chromosomal instability. *Journal of the National Cancer Institute*, **105**, 1202-1211.

Shay, J.W. & Wright, W.E. (2011) Role of telomeres and telomerase in cancer. *Seminars in cancer biology*, **21**, 349-353.

Sieglova, Z., Zilovcova, S., Cermak, J., Rihova, H., Brezinova, D., Dvorakova, R., Markova, M., Maaloufova, J., Sajdova, J., Brezinova, J., Zemanova, Z. & Michalova, K. (2004) Dynamics of telomere erosion and its association with genome instability in myelodysplastic syndromes (MDS) and acute myelogenous leukemia arising from MDS: a marker of disease prognosis? *Leukemia research*, **28**, 1013-1021.

Tothova, Z., Steensma, D.P. & Ebert, B.L. (2013) New Strategies in Myelodysplastic Syndromes: Application of molecular diagnostics to clinical practice. *Clinical cancer research : an official journal of the American Association for Cancer Research*.

Townsley, D.M., Dumitriu, B. & Young, N.S. (2014) Bone marrow failure and the telomeropathies. *Blood*, **124**, 2775-2783.

Vardiman, J.W., Thiele, J., Arber, D.A., Brunning, R.D., Borowitz, M.J., Porwit, A., Harris, N.L., Le Beau, M.M., Hellstrom-Lindberg, E., Tefferi, A. & Bloomfield, C.D. (2009) The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: rationale and important changes. *Blood*, **114**, 937-951.

Telomere length is prognostic in MDS but not AML

- 464 Walter, M.J., Shen, D., Ding, L., Shao, J., Koboldt, D.C., Chen, K., Larson, D.E., McLellan, M.D., Dooling,  
465 D., Abbott, R., Fulton, R., Magrini, V., Schmidt, H., Kalicki-Veizer, J., O'Laughlin, M., Fan, X.,  
466 Grillo, M., Witowski, S., Heath, S., Frater, J.L., Eades, W., Tomasson, M., Westervelt, P.,  
467 DiPersio, J.F., Link, D.C., Mardis, E.R., Ley, T.J., Wilson, R.K. & Graubert, T.A. (2012) Clonal  
468 architecture of secondary acute myeloid leukemia. *The New England journal of medicine*,  
469 **366**, 1090-1098.
- 470 Xie, M., Lu, C., Wang, J., McLellan, M.D., Johnson, K.J., Wendl, M.C., McMichael, J.F., Schmidt, H.K.,  
471 Yellapantula, V., Miller, C.A., Ozenberger, B.A., Welch, J.S., Link, D.C., Walter, M.J., Mardis,  
472 E.R., DiPersio, J.F., Chen, F., Wilson, R.K., Ley, T.J. & Ding, L. (2014) Age-related mutations  
473 associated with clonal hematopoietic expansion and malignancies. *Nature medicine*, **20**,  
474 1472-1478.
- 475 Yamaguchi, H., Baerlocher, G.M., Lansdorp, P.M., Chanock, S.J., Nunez, O., Sloand, E. & Young, N.S.  
476 (2003) Mutations of the human telomerase RNA gene (TERC) in aplastic anemia and  
477 myelodysplastic syndrome. *Blood*, **102**, 916-918.
- 478 Young, N.S. (2010) Telomere biology and telomere diseases: implications for practice and research.  
479 *Hematology / the Education Program of the American Society of Hematology. American*  
480 *Society of Hematology. Education Program*, **2010**, 30-35.
- 481
- 482
- 483
- 484
- 485
- 486
- 487
- 488
- 489
- 490
- 491
- 492
- 493
- 494
- 495
- 496
- 497
- 498
- 499

Telomere length is prognostic in MDS but not AML

Figures and Legends.

Table 1. Demographics of MDS and AML Cohorts

**Figure 1.** (A) Telomere length in the MDS cohort was significantly longer than those of the AML cohort despite being older median age 68 v 56 years ( $p < .0001$ ). B) In MDS patients there was no correlation between telomere length and age of diagnosis ( $\rho^2 = .0212$ ;  $P = .2$ ). (C) In AML patients there was a positive correlation between telomere length and age of diagnosis ( $\rho^2 = .0890$ ;  $P = .003$ ). (D) Older AML patients (age  $>60$ ) had significantly longer telomeres than younger patients ( $P = .02$ ).

**Figure 2.** (A) In the MDS cohort there was a significant association between the telomere length and the number of cytopenia but not with (B) cytogenetic risk group ( $P = .6$  for trend) or (C) IPSS sub-groups ( $\rho^2 = .14$  for correlation) or (D) blast counts ( $\rho^2 = .22$   $P = .503$ ). (E) Patients with a FLT3-ITD showed significantly shorter telomeres than the FLT3-WT group (overall  $P = .03$  and there was a trend towards shorter telomere length in the FLT3-ITD group when compared with the FLT3-TKD group ( $P = .12$ )). F) Telomerase activity was significantly higher in AML CD34 + cells compared to MDS CD34+ cells ( $P = .0002$ ).

**Figure 3.** (A) Using our previously described CLL fusogenic length threshold ( $TL \leq 3.81$  kb) there is no difference in survival in AML patients ( $HR = 1.47$  (0.80-2.68),  $P = 0.2$ ). (B) In contrast, categorization of the MDS cohort above and below the fusogenic length threshold (3.81 kb) demonstrated that MDS patients with short telomeres had an inferior survival ( $HR = 5.0$  (2.7-10.0),  $P < .0001$ ).

**Supplementary Figure 1.** (A) In 20 MDS patients bone marrow mononuclear cells were sorted but there was no significant difference in telomere length in CD34+ and CD34 – selected sub-populations. (B) In the MDS cohort longer telomere length was associated with female sex ( $P = .01$ ).



Telomere length is prognostic in MDS but not AML

**Supplementary Figure 2.** (A) In the MDS cohort there was a significant association between the number of cytopenias and overall survival ( $P < .0001$ ). (B) In the MDS cohort patients with high risk cytogenetic lesions tended to have worse survival although this did not reach statistical significance probably due to the relatively small sample size ( $P = .12$ ).

**Supplementary Figure 3.** Quartile analysis of telomere length revealed no significant difference in (A) overall survival and (B) relapse-free survival in the de novo AML cohort. In contrast, (C) MDS patients in the lower two telomere length quartiles showed significantly shorter overall survival.

For Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Unable to Convert Image

The dimensions of this image (in pixels) are too large to be converted. For this image to convert, the total number of pixels (height x width) must be less than 40,000,000 (40 megapixels).

Or Peer Review




## Unable to Convert Image

The dimensions of this image (in pixels) are too large to be converted. For this image to convert, the total number of pixels (height x width) must be less than 40,000,000 (40 megapixels).

Or Peer Review

Demographic Feature	MDS Cohort (n=80)	AML Cohort (n=95)
Sex		
Male	50	48
Female	30	47
Age		
Median (range)	68 (21-86)	56 (17-80)
<60	20	56
≥60	60	39
Cytogenetic Risk Group		
Good	45	6
Intermediate	15	76
Poor	15	3
Failed/Not known	5	10
Number of cytopenias		
1	37	
2	15	
3	28	
Bone Marrow Blast Count		
<5%	36	
5-10%	13	
10-20%	24	
>20%	7	
IPSS Score		
High	25	
Intermediate-2	11	
Intermediate-1	19	
Low	25	
Treatment		
Intensive		89
Non-intensive		6
ITD		
WT		49
Mutant		39
Unknown		7
NPM1c		
WT		35
Mutant		54
Unknown		6
TKD		
WT		73
Mutant		19
Unknown		3
WBC		
Median (range)		34.1 (1.3-294.0)
Performance Status (WHO)		
0		54
1		30
2		4
3+		7


Williams et al\_Supplementary Figure 1



Supplementary Figure 1

177x283mm (600 x 600 DPI)


Williams et al\_Supplementary Figure 2



Supplementary Figure 2

165x244mm (600 x 600 DPI)

Williams et al\_Supplementary Figure 3



Supplementary Figure 3

226x416mm (600 x 600 DPI)